

Low-lying fermion modes of $N_f = 2$ improved Wilson fermions*

QCDSF and UKQCD Collaboration: R. Horsley^a, T.G. Kovács^a, V. Linke^b, D. Pleiter^a, G. Schierholz^{a,c}

^aJohn von Neumann Institute NIC / DESY Zeuthen, D-15738 Zeuthen, Germany

^bInstitut für Theoretische Physik, Freie Universität Berlin, D-14195 Berlin

^cDeutsches Elektronen-Synchrotron DESY, D-22603 Hamburg, Germany

We present preliminary results for the topological charge and susceptibility determined from the low-lying eigenmodes of the Wilson-Dirac operator. These modes have been computed on dynamical configurations with $N_f = 2$ non-perturbatively improved Wilson fermions. We compare our results with the eigenmodes of fermions in the quenched approximation.

1. INTRODUCTION

For several reasons there is particular interest in the low-lying eigenmodes of the Dirac operator in QCD. From phenomenological models it is expected that for sufficiently light quark masses physics, e.g. hadron correlators, is dominated by these modes. Calculations on the lattice enable us to study the relevance of low-lying eigenmodes for various observables directly.

Furthermore, the eigenmodes of the Dirac operator carry information about the topological content of the background gauge field. The topological properties can also be probed using gluonic methods. These however typically suffer from large fluctuations on very short scales and therefore require some sort of smoothing procedures. Usually the quantities of interest are affected by the applied filtering method.

Results from fermionic methods on the other hand are expected to be sensitive to the effects caused by explicitly breaking chiral symmetry when using Wilson fermions. Although the symmetry is expected to be restored in the continuum limit, these effects are potentially large on coarse lattices. However, these can be reduced by applying Symanzik's improvement program. Removing discretization effects of $O(a)$, where a is the lattice spacing, is achieved by adding a single counterterm to the action. The improved Wilson-

Dirac operator then reads:

$$M = 1 - \kappa \left[H + \frac{i}{2} c_{\text{SW}} F_{\mu\nu} \sigma_{\mu\nu} \right], \quad (1)$$

where H is the hopping term of the usual Wilson fermion action.

For the calculations presented here we used $O(20-50)$ dynamical configurations with $N_f = 2$ flavours of degenerate quarks for six different combinations of β and κ^{sea} . We used lattices of size $16^3 \times 32$ with a lattice spacing a that varies between 0.104(1) and 0.095(1) fm (using the force scale $r_0 = 0.5$ fm). For the lightest and heaviest sea quarks the ratio $m_{\text{PS}}/m_{\text{V}}$ was 0.60 and 0.83, respectively. For comparison we performed the same calculations using quenched configurations at $\beta = 6.0$ and $a = 0.093(1)$ fm.

2. COMPUTING LOW-LYING EIGENVALUES

The Wilson-Dirac operator M is a non-hermitian operator. The chirality ω_i of an eigenmode λ_i can be calculated using the corresponding right eigenvector v_i :

$$\omega_i = \sum_t \omega_i(t) = \sum_t \sum_x v_i^\dagger(x, t) \gamma_5 v_i(x, t). \quad (2)$$

We can distinguish between two types of eigenmodes. Firstly, complex modes with non-vanishing imaginary part and $\omega_i = 0$ and, secondly, real modes with vanishing imaginary part and $\omega_i \neq 0$.

*Talk given by D. Pleiter at the Lattice Conference 2001, Berlin, Germany.




Figure 1. $O(100)$ smallest eigenvalues of the massless Dirac operator calculated on 20 gauge dynamical (left) and quenched (right) configurations.

In principle, it is possible to find the real modes by determining the zero modes of the hermitian operator $\gamma_5 M$. If $\lambda = 1 - \kappa\sigma$ is a real mode of M , then $\gamma_5 M$ will have a zero mode for $\kappa = 1/\sigma$. In practice it might however be difficult to detect several real modes which lie close to each other. We therefore decided to calculate the 70-100 smallest eigenvalues of the non-hermitian operator M using the Arnoldi algorithm [1]. Results for the massless Dirac operator

$$D = \frac{1}{2\kappa} M - \frac{1}{2} \left(\frac{1}{\kappa_c} - \frac{1}{\kappa} \right) \quad (3)$$

are plotted in Fig. 1.

3. REAL MODES

For dynamical fermions one would expect eigenmodes around zero to be suppressed. As can be seen from Fig. 1, this seems indeed to be the case. This effect is more pronounced if one looks at the density of the real modes (Fig. 2).

Real modes close to (or below) zero, which correspond to zero modes of $\gamma_5 M$ at $\kappa \lesssim \kappa_c$, are believed to be the origin of so-called exceptional configurations, which are encountered frequently in the quenched approximation when approaching




Figure 2. Density of the real modes as a function of the real part. The area under the curve is normalized to one. The graphs are ordered from heaviest, i.e. infinite, (top) to smallest (bottom) sea quark mass.

the chiral regime. Although these modes are expected to be suppressed for dynamical fermions, there is a non-zero probability for this kind of configurations to occur (see Fig. 2).

The correlation of the chirality $c_i(r) = (1/V) \sum_{x,t} \omega_i(x,t) \omega_i(x+r,t)$ shown in Fig. 3 was found to be narrow, confirming the observations made in [2]. Typically the chirality of small real modes is concentrated in a few time slices.

4. TOPOLOGICAL SUSCEPTIBILITY

The topological susceptibility normalised by the volume is defined by

$$\chi = \frac{\langle Q_{\text{top}}^2 \rangle}{V}. \quad (4)$$

The topological charge Q_{top} can be obtained via the Atiyah-Singer index theorem, which is approximately true on the lattice. It establishes a relation between the difference in the number of real modes with negative (n^+) and positive (n^-) chirality and the topological charge:

$$Q_{\text{top}} = n^+ - n^- \quad (5)$$

For a subset of configurations we compared the results for Q_{top} with those obtained from gluonic




Figure 3. Chirality per timeslice $\omega_i(t)$ for real modes (upper picture) and the normalized correlation function of the chirality (lower picture) $c_i(r)$ on a configuration at $\beta = 5.29$, $\kappa^{\text{sea}} = 0.13450$. The real mode with the smallest eigenvalue is plotted with a solid line.

operators using the Boulder smoothing technique. The maximum difference found was ± 2 . For a more detailed comparison see [3].

The results for the topological susceptibility are plotted in Fig. 4. The discrepancy between the results from the fermionic method applied in this work and from the gluonic method used by Hart and Teper is likely to be explained by a lack of statistics. Another source of uncertainty arises from the fact that we only calculated the lowest 70-100 eigenvalues. Although the probability for real modes is expected to decrease when moving away from zero, it does not become zero. To check for this possible error, we looked at the topological susceptibility as a function of the largest real mode taken into account (see Fig. 5). For all data sets we found a reasonable plateau.

5. CONCLUSIONS

Comparing real modes of quenched and dynamical configurations we found evidence for an expected depopulation of the region around zero. The results for Q_{top} obtained by using the Atiya-Singer-theorem were in reasonable agreement with numbers obtained from gluonic meth-




Figure 4. Topological susceptibility as a function of the pseudoscalar meson mass. The solid symbols show the results from this work, while UKQCD results using gluonic methods [4] are plotted with open symbols.




Figure 5. Topological susceptibility versus the cut-off of used real modes.

ods. Our preliminary results for the topological susceptibility were found to be similar to other calculations using the same configurations but gluonic methods.

ACKNOWLEDGEMENTS

The numerical calculations were performed on the Hitachi SR8000 at LRZ (Munich), the Cray T3E at EPCC (Edinburgh) and ZIB (Berlin). We wish to thank all institutions for their support.

REFERENCES

1. R.B. Lehoucq, D.C. Sorensen, C. Yang, "ARPACK Users Guide", 1997; K. Maschhoff, D. Sorensen, 1996.
2. T. DeGrand, A. Hasenfratz, Phys. Rev. D **64** (2001) 034512.
3. A. Hart, this conference.
4. A. Hart, M. Teper [hep-lat/0108022].